

# Lamin B1 Rabbit pAb

Catalog No.: A1910 **48 Publications**

## Basic Information

### Observed MW

68kDa/45kDa

### Calculated MW

66kDa

### Category

Primary antibody

### Applications

WB, IHC-P, IF/ICC, IP, ELISA, ChIP

### Cross-Reactivity

Human, Mouse, Rat

## Background

This gene encodes one of the two B-type lamin proteins and is a component of the nuclear lamina. A duplication of this gene is associated with autosomal dominant adult-onset leukodystrophy (ADLD). Alternative splicing results in multiple transcript variants.

## Recommended Dilutions

**WB** 1:500 - 1:1000

**IHC-P** 1:50 - 1:200

**IF/ICC** 1:50 - 1:200

**IP** 0.5µg-4µg antibody for  
200µg-400µg extracts of  
whole cells

**ELISA** Recommended starting  
concentration is 1 µg/mL.  
Please optimize the  
concentration based on  
your specific assay  
requirements.

**ChIP** 5µg antibody for  
10µg-15µg of Chromatin

## Immunogen Information

### Gene ID

4001

### Swiss Prot

P20700

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 397-586 of human Lamin B1 (NP\_005564.1).

### Synonyms

LMN; ADLD; LMN2; LMNB; MCPH26; Lamin B1

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 50% glycerol, pH7.3.

## Contact

---


 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)


 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

---


## Validation Data



Western blot analysis of various lysates using Lamin B1 Rabbit pAb (A1910) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1min.




Immunoprecipitation of Lamin B1 in 300 µg extracts from Jurkat cells using 0.5 µg Lamin B1 Rabbit pAb (A1910). Western blot analysis was performed using Lamin B1 Rabbit pAb (A1910) at 1:1000 dilution.




Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg Lamin B1 antibody (A1910). Western blot was performed from the immunoprecipitate using Lamin B1 (A1910) at a dilution of 1:1000.


## Validation Data



Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg Lamin B1 antibody (A1910). Western blot was performed from the immunoprecipitate using Lamin B1 antibody (A1910) at a dilution of 1:1000.




Western blot analysis of various lysates using Lamin B1 Rabbit pAb (A1910) at 1:1000 dilution. Jurkat cells were treated by Etoposide (25 µM) at 37°C for 5 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.




Western blot analysis of various lysates using Lamin B1 Rabbit pAb (A1910) at 1:1000 dilution. L929 cells were treated by staurosporine(1 µM) for 3 hour. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.


## Validation Data




Chromatin immunoprecipitation was performed with cross-linked chromatin from HeLa, using Lamin-B1 Rabbit mAb (A1910) and rabbit IgG(AC042). The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram compares the ratio of the immunoprecipitated DNA versus the input.




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using Lamin B1 Rabbit pAb antibody (A1910) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.






Confocal immunofluorescence analysis of HeLa cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:200. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of C6 cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of NIH/3T3 cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



## Validation Data


---

Confocal immunofluorescence analysis of HeLa cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:200. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of paraffin-embedded Human breast cancer using Lamin B1 Rabbit pAb (A1910) at dilution of 1:150 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.

Immunofluorescence analysis of PC-12 cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded Mouse kidney using Lamin B1 Rabbit pAb (A1910) at dilution of 1:150 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.

Immunofluorescence analysis of U2OS cells using Lamin B1 Rabbit pAb (A1910) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain using Lamin B1 Rabbit pAb (A1910) at dilution of 1:150 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.