

SHP1 Rabbit mAb

Catalog No.: A19111 **Recombinant** **1 Publications**

Basic Information

Observed MW

68kDa

Calculated MW

68kDa

Category

Primary antibody

Applications

ELISA,WB,IHC-P,IF/ICC

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC0295

Background

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. N-terminal part of this PTP contains two tandem Src homolog (SH2) domains, which act as protein phospho-tyrosine binding domains, and mediate the interaction of this PTP with its substrates. This PTP is expressed primarily in hematopoietic cells, and functions as an important regulator of multiple signaling pathways in hematopoietic cells. This PTP has been shown to interact with, and dephosphorylate a wide spectrum of phospho-proteins involved in hematopoietic cell signaling. Multiple alternatively spliced variants of this gene, which encode distinct isoforms, have been reported.

Recommended Dilutions

WB	1:500 - 1:2000
IHC-P	1:200 - 1:800
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

5777

Swiss Prot

P29350

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 496-595 of human SHP1 (P29350).

Synonyms

HCP; HCPH; SHP1; SHP-1; HPTP1C; PTP-1C; SHP-1L; SH-PTP1

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.


Validation Data




Western blot analysis of extracts of Raji cells, using SHP1 antibody (A19111) at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.




Western blot analysis of extracts of various cell lines, using SHP1 antibody (A19111) at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 3min.



Immunofluorescence analysis of rat spleen using SHP1 Rabbit mAb (A19111) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




Immunofluorescence analysis of mouse spleen using SHP1 Rabbit mAb (A19111) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of SHP1 in paraffin-embedded human spleen tissue using SHP1 Rabbit mAb (A19111) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.


Validation Data



Immunohistochemistry analysis of SHP1 in paraffin-embedded human tonsil tissue using SHP1 Rabbit mAb (A19111) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of SHP1 in paraffin-embedded mouse spleen tissue using SHP1 Rabbit mAb (A19111) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of SHP1 in paraffin-embedded rat spleen tissue using SHP1 Rabbit mAb (A19111) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.