

SCAF8 Rabbit pAb

Catalog No.: A19467

Basic Information

Observed MW

170kDa

Calculated MW

141kDa

Category

Primary antibody

Applications

ELISA, WB

Cross-Reactivity

Human, Mouse, Rat

Background

Enables RNA binding activity and RNA polymerase II C-terminal domain phosphoserine binding activity. Involved in negative regulation of termination of RNA polymerase II transcription, poly(A)-coupled and positive regulation of DNA-templated transcription, elongation. Located in nucleoplasm.

Recommended Dilutions

WB 1:500 - 1:1000

Immunogen Information

Gene ID

22828

Swiss Prot

Q9UPN6

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 572-634 of human SCAF8 (NP_055707.3).

Synonyms

RBM16; SCAF8

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts of various cell lines, using SCAF8 antibody (A19467) at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 μ g per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.



Western blot analysis of extracts of HeLa cells, using SCAF8 antibody (A19467) at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 μ g per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 180s.