## NF-κB2 Rabbit mAb

Catalog No.: A19605 Recombinant 2 Publications



## **Basic Information**

### **Observed MW**

120kDa

#### **Calculated MW**

97kDa

### Category

Primary antibody

### **Applications**

WB,IHC-P,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

#### CloneNo number

ARC0084

## **Background**

This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protein encoded by this gene can function as both a transcriptional activator or repressor depending on its dimerization partner. The p100 full-length protein is co-translationally processed into a p52 active form. Chromosomal rearrangements and translocations of this locus have been observed in B cell lymphomas, some of which may result in the formation of fusion proteins. There is a pseudogene for this gene on chromosome 18. Alternative splicing results in multiple transcript variants.

# **Recommended Dilutions**

**WB** 1:1000 - 1:6000

IHC-P 1:100 - 1:500

**ELISA** Recommended starting concentration is 1 μg/mL.

Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

**Gene ID Swiss Prot** 4791 Q00653

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 745-877 of human NF-κB2 (Q00653).

## **Synonyms**

p52; p100; H2TF1; LYT10; CVID10; LYT-10; NF-kB2; p49/p100; NF-kB2

## **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\overline{\triangle}$	ī	www.ahclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.




Western blot analysis of various lysates using NF- $\kappa$ B2 Rabbit mAb (A19605) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 $\mu$ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3s.



Immunohistochemistry analysis of paraffinembedded Human appendix using NF-κB2 Rabbit mAb (A19605) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.