# PGC1α/β Rabbit mAb

Catalog No.: A19674 Recombinant 13 Publications



# **Basic Information**

### **Observed MW**

113kDa

#### **Calculated MW**

113kDa

### Category

Primary antibody

### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Mouse, Rat

#### CloneNo number

ARC0153

# **Background**

The protein encoded by this gene is a transcriptional coactivator that regulates the genes involved in energy metabolism. This protein interacts with PPARgamma, which permits the interaction of this protein with multiple transcription factors. This protein can interact with, and regulate the activities of, cAMP response element binding protein (CREB) and nuclear respiratory factors (NRFs). It provides a direct link between external physiological stimuli and the regulation of mitochondrial biogenesis, and is a major factor that regulates muscle fiber type determination. This protein may be also involved in controlling blood pressure, regulating cellular cholesterol homoeostasis, and the development of obesity.

# **Recommended Dilutions**

**WB** 1:1000 - 1:6000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# Immunogen Information

 Gene ID
 Swiss Prot

 10891/133522
 Q9UBK2/Q86YN6

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

LEM6; PGC-1(alpha); PGC-1alpha; PGC-1v; PGC1; PGC1A; PPARGC1; ERRL1; PERC; PGC-1(beta); PGC1B; PGC1 $\alpha/\beta$ 

# **Contact**

<b>a</b>	400-999-6126
$\sim$	cn.market@abclonal.com.cn
<b>⊙</b>	www.abclonal.com.cn

# **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

# **Validation Data**



Western blot analysis of various lysates using PGC1 $\alpha$ / $\beta$  Rabbit mAb (A19674) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 $\mu$ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.