

# FANCD2 Rabbit mAb

Catalog No.: A19692 **Recombinant** **1 Publications**

## Basic Information

### Observed MW

166kDa

### Calculated MW

164kDa

### Category

Primary antibody

### Applications

ELISA,WB,IP

### Cross-Reactivity

Human, Mouse, Rat

### CloneNo number

ARC0172

## Background

The Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCI, FANCI (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The previously defined group FANCI is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity; they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group D2. This protein is monoubiquitinated in response to DNA damage, resulting in its localization to nuclear foci with other proteins (BRCA1 AND BRCA2) involved in homology-directed DNA repair. Alternative splicing results in multiple transcript variants.

## Recommended Dilutions

**WB** 1:500 - 1:1000

**IP** 0.5µg-4µg antibody for  
200µg-400µg extracts of  
whole cells

## Immunogen Information

### Gene ID

2177

### Swiss Prot

Q9BXW9

### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 200-300 of human FANCD2 (Q9BXW9).

### Synonyms

FA4; FAD; FADC; FAD2; FA-D2; FANCD; FANCD2

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.


## Validation Data



Immunoprecipitation analysis of 300ug extracts of HeLa cells using 3ug FANCD2 Rabbit mAb (A19692). Western blot was performed from the immunoprecipitate using FANCD2 Rabbit mAb (A19692) at a dilution of 1:1000.



Western blot analysis of various lysates using FANCD2 Rabbit mAb (A19692) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.



Western blot analysis of lysates from mouse testis, using FANCD2 Rabbit mAb (A19692) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 3min.