

PRK2/PKN2 Rabbit mAb

Catalog No.: A19746 **Recombinant** **1 Publications**

Basic Information

Observed MW

112kDa

Calculated MW

112kDa

Category

Primary antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC2274

Background

Enables RNA polymerase binding activity; histone deacetylase binding activity; and protein serine/threonine kinase activity. Involved in several processes, including apical junction assembly; positive regulation of cell cycle; and positive regulation of viral genome replication. Located in several cellular components, including cleavage furrow; cytoskeleton; and midbody.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

5586

Swiss Prot

Q16513

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

PAK2; PRK2; STK7; Pak-2; PRKCL2; PRO2042; PRK2/PKN2

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of various lysates using PRK2/PRK2/PKN2 Rabbit mAb (A19746) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.