

# [KO Validated] STRAP Rabbit pAb

Catalog No.: A19989 **KO** Validated

## Basic Information

### Observed MW

39kDa

### Calculated MW

38kDa

### Category

Primary antibody

### Applications

WB, ELISA

### Cross-Reactivity

Human, Mouse, Rat

## Background

Enables RNA binding activity. Involved in maintenance of gastrointestinal epithelium; negative regulation of transforming growth factor beta receptor signaling pathway; and spliceosomal snRNP assembly. Located in cytosol. Part of SMN complex. Implicated in adenocarcinoma; colorectal carcinoma; large cell carcinoma; lung carcinoma; and squamous cell neoplasm.

## Recommended Dilutions

**WB** 1:1000 - 1:2000

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

11171

### Swiss Prot

Q9Y3F4

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

MAWD; PT-WD; UNRIP; AP

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

## Validation Data

---



Western blot analysis of lysates from wild type (WT) and STRAP knockout (KO) 293T cells, using [KO Validated] STRAP Rabbit pAb (A19989) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 $\mu$ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.