# **β-Catenin Mouse mAb**

Catalog No.: A20221 3 Publications



# **Basic Information**

#### **Observed MW**

92kDa

### **Calculated MW**

85kDa

### Category

Primary antibody

### **Applications**

WB,IHC-P,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

#### CloneNo number

AMC0440

# **Background**

The protein encoded by this gene is part of a complex of proteins that constitute adherens junctions (AJs). AJs are necessary for the creation and maintenance of epithelial cell layers by regulating cell growth and adhesion between cells. The encoded protein also anchors the actin cytoskeleton and may be responsible for transmitting the contact inhibition signal that causes cells to stop dividing once the epithelial sheet is complete. Finally, this protein binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatrixoma (PTR), medulloblastoma (MDB), and ovarian cancer. Alternative splicing results in multiple transcript variants.

# **Recommended Dilutions**

**WB** 1:1000 - 1:4000

IHC-P 1:100 - 1:500

**ELISA** Recommended starting concentration is 1 μg/mL.

Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

 Gene ID
 Swiss Prot

 1499
 P35222

### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 300-400 of human  $\beta$ -Catenin (NP\_001895.1).

### **Synonyms**

EVR7; CTNNB; MRD19; NEDSDV; armadillo; β-Catenin

## **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	T	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationMouseIgG1,KappaAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.




Western blot analysis of various lysates using  $\beta$ -Catenin Mouse mAb (A20221) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.


Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Immunohistochemistry analysis of paraffinembedded Human liver cancer using  $\beta$ -Catenin Mouse mAb (A20221) at dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human skin using  $\beta$ -Catenin Mouse mAb (A20221) at dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.