

# OsDLT Rabbit pAb

**Catalog No.: A20318**

## Basic Information

### Observed MW

66kDa

### Calculated MW

66kDa

### Category

Primary antibody

### Applications

ELISA, WB

### Cross-Reactivity

Oryza sativa

## Background

Probable transcription factor that acts as positive regulator of brassinosteroid (BR) signaling. Functions downstream of BRI1 and GSK2 to modulate BR responses. Acts as a direct target of GSK2 kinase to mediate BR responses. Involved in feedback inhibition of BR biosynthetic genes. Repressed by BZR1. Cooperatively functions in a transactivating complex with SMOS1 to enhance the transcription of the SMOS1 target PH1-1, and regulate plant organ size. Interaction between SMOS1 and DLT is a crosstalk point for auxin and brassinosteroid signaling

## Recommended Dilutions

**WB** 1:500 - 1:2000

## Immunogen Information

### Gene ID

4339978

### Swiss Prot

Q9LWU9

### Immunogen

Recombinant protein of Oryza sativa OsDLT.

### Synonyms

DLT; OsGRAS-32; D62; GS6; SMOS2; OsDLT

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

# Validation Data



Western blot analysis of extracts of various tissues from the japonica rice (*Oryza sativa* L.) variety Zhonghua 11, using OsDLT Rabbit pAb (A20318) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Enhanced Kit (RM00021).  
Exposure time: 30s.