# AP1G1 Rabbit pAb

Catalog No.: A20454



# **Basic Information**

#### **Observed MW**

91kDa

#### **Calculated MW**

91kDa

### Category

Primary antibody

### **Applications**

ELISA,WB,IF/ICC

### **Cross-Reactivity**

Human, Mouse

# **Background**

Adaptins are important components of clathrin-coated vesicles transporting ligand-receptor complexes from the plasma membrane or from the trans-Golgi network to lysosomes. The adaptin family of proteins is composed of four classes of molecules named alpha, beta-, beta prime- and gamma- adaptins. Adaptins, together with medium and small subunits, form a heterotetrameric complex called an adaptor, whose role is to promote the formation of clathrin-coated pits and vesicles. The protein encoded by this gene is a gamma-adaptin protein and it belongs to the adaptor complexes large subunits family. Two transcript variants encoding different isoforms have been found for this gene.

## **Recommended Dilutions**

**WB** 1:500 - 1:1000

**IF/ICC** 1:50 - 1:200

# **Immunogen Information**

**Gene ID Swiss Prot** 164 043747

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-240 of human AP1G1 (NP\_001025178).

### **Synonyms**

ADTG; CLAPG1; USRISD; AP1G1

### **Contact**

| <b>a</b>       | 400-999-6126              |
|----------------|---------------------------|
| <b>×</b>       | cn.market@abclonal.com.cn |
| $\overline{a}$ | www.abclonal.com.cn       |


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.




Western blot analysis of various lysates using AP1G1 Rabbit pAb (A20454) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 20s.



Immunofluorescence analysis of NIH/3T3 cells using AP1G1 Rabbit pAb (A20454) at dilution of 1:200 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.