

# IL1 $\beta$ Rabbit pAb

Catalog No.: A20529 **7 Publications**

## Basic Information

### Observed MW

31kDa

### Calculated MW

31kDa

### Category

Primary antibody

### Applications

WB,ELISA

### Cross-Reactivity

Human, Mouse, Rat

## Background

Enables cytokine activity. Involved in several processes, including positive regulation of intracellular signal transduction; regulation of gene expression; and regulation of neurogenesis. Located in extracellular space. Used to study several diseases, including artery disease (multiple); brain disease (multiple); glomerulonephritis (multiple); interstitial lung disease (multiple); and kidney failure (multiple). Biomarker of several diseases, including artery disease (multiple); auditory system disease (multiple); brain disease (multiple); kidney failure (multiple); and lung disease (multiple). Human ortholog(s) of this gene implicated in several diseases, including autoimmune disease (multiple); blood platelet disease (multiple); eye disease (multiple); gastrointestinal system cancer (multiple); and lung disease (multiple). Orthologous to human IL1B (interleukin 1 beta).

## Recommended Dilutions

**WB** 1:500 - 1:1000

**ELISA** Recommended starting concentration is 1  $\mu$ g/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

24494

### Swiss Prot

Q63264

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 117-268 of rat IL1 $\beta$  (NP\_113700.2).

### Synonyms

IL-1F2; IL1 $\beta$

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of various lysates using IL1 $\beta$  Rabbit pAb (A20529) at 1:1000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25  $\mu$ g per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Enhanced Kit (RM00021).  
Exposure time: 180s.



Western blot analysis of lysates from THP-1 cells using IL1 $\beta$  Rabbit pAb (A20529) at 1:1000 dilution. THP-1 cells were treated by LPS (1  $\mu$ g/ml) at 37°C for 8 hours.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25  $\mu$ g per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 90s.