

# Ghd 7.1 Rabbit pAb

**Catalog No.: A20584**

## Basic Information

**Observed MW**

80kDa

**Calculated MW**

80kDa

**Category**

Primary antibody

**Applications**

WB, ELISA

**Cross-Reactivity**

Oryza sativa

## Background

Probable transcription factor involved in the regulation of flowering time under long day (LD) conditions. Functions as repressor of flowering. Controls flowering time by negatively regulating the expression of HD3A. Acts downstream of the phytochrome B to repress the expression of EHD1, an activator of the flowering promoter genes HD3A and RFT1. Controls photoperiodic flowering response. Seems to be one of the component of the circadian clock. Expression of several members of the ARR-like family is controlled by circadian rhythm. The particular coordinated sequential expression of PRR73, PRR37, PRR95, PRR59 and PPR1 result to circadian waves that may be at the basis of the endogenous circadian clock (By similarity).

## Recommended Dilutions

**WB** 1:500 - 1:2000**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

**Gene ID**

4344399

**Swiss Prot**

Q0D3B6

**Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

**Synonyms**

HD2; DTH7; PRR37; OsPRR37; Ghd 7.1

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**

Affinity purification


**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

## Validation Data

---



Western blot analysis of extracts of various tissues from the japonica rice (*Oryza sativa* L.) variety Zhonghua 11, using Ghd 7.1 Rabbit pAb (A20584) at 1:1000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Enhanced Kit (RM00021).  
Exposure time: 60s.