

CD70 Rabbit mAb

Catalog No.: A20589 **Recombinant**

Basic Information

Observed MW

21-27kDa

Calculated MW

21kDa

Category

Primary antibody

Applications

WB,FC,ELISA

Cross-Reactivity

Human

CloneNo number

ARC5081-01

Background

The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis.

Recommended Dilutions

WB 1:500 - 1:2000**FC** 1:100 - 1:800

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

970

Swiss Prot

P32970

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CD27L; LPFS3; CD27-L; CD27LG; TNFSF7; TNLG8A; CD70

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data




Western blot analysis of various lysates using CD70 Rabbit mAb (A20589) at dilution incubated overnight at 4°C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.



Flow cytometry: 1×10^6 Raji cells were stained with CD70 Rabbit mAb (A20589, 1 µg/ml Blue line). Non-fluorescently stained Raji cells were used as blank control (red line).