

HCoV-NL63 Spike S2 Rabbit pAb

Catalog No.: A20611

Basic Information

Observed MW

160kDa

Calculated MW

150kDa

Category

Primary antibody

Applications

ELISA, WB

Cross-Reactivity

HCoV-NL63

Background

S1 region attaches the virion to the cell membrane by interacting with host ACE2, initiating the infection. Binding to the receptor probably induces conformational changes in the S glycoprotein unmasking the fusion peptide and activating membranes fusion. S2 region belongs to the class I viral fusion protein. Under the current model, the protein has at least 3 conformational states: pre-fusion native state, pre-hairpin intermediate state, and post-fusion hairpin state. During viral and target cell membrane fusion, the coiled coil regions (heptad repeats regions assume a trimer-of-hairpins structure, positioning the fusion peptide in close proximity to the C-terminal region of the ectodomain. The formation of this structure appears to drive apposition and subsequent fusion of viral and target cell membranes.

Recommended Dilutions

WB 1:500 - 1:1000

Immunogen Information

Gene ID

2943499

Swiss Prot

Q6Q1S2

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1200-1300 of coronavirus Spike S2 (YP_003767.1).

Synonyms

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from Human coronavirus (HCoV-NL63) Spike Protein (S1+S2 ECDHis Tag), using HCoV-NL63 Spike S2 Rabbit pAb (A20611) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.