Leader in Biomolecular Solutions for Life Science

OsFD2 Rabbit pAb

Catalog No.: A20657



Basic Information

Observed MW 18kDa

Calculated MW 18kDa

Category Primary antibody

Applications WB,ELISA

Cross-Reactivity Oryza sativa

Background

Overexpression of OsFD2, a member of the Poaceae FD2 group, produced smaller leaves with shorter plastochrons, suggesting that OsFD2 controls leaf development.

Recommended Dilutions

WB	1:500 - 1:2000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID 9266154 Swiss Prot Q5Z4G3

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms OsFD2; OsbZIP55

Contact


6	400-999-6126
\times	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

Product Information

Source Rabbit **Isotype** IgG Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of extracts of various tissues from the japonica rice (Oryza sativa L.) variety Zhonghua 11, using OsFD2 Rabbit pAb (A20657) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per Iane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 600s.