2'-O-Methylguanosine(Gm) Rabbit pAb

www.abclonal.com

ABclonal

Catalog No.: A20697

Basic Information

Observed MW

Refer to figures

Calculated MW

Category

Primary antibody

Applications

DB,ELISA

Cross-Reactivity

Species independent

Background

RNA methylation plays a significant regulatory role in various of physiological activities and it has gradually become a hotspot of epigenetics in the past decade. 2'-O-methyladenosine (Am), 2'-O-methylguanosine (Gm),2'-O-methylcytidine (Cm),2'-O-methyluridine (Um),N 6-methyladenosine (m6A), N 1-methylguanosine (m1G),5-methylcytidine (m5C),and 5-methyluridine (m5U) are representative 2'-O-methylation and base-methylation modified epigenetic marks of RNA. 2'-O-methyltransferase is a modified nucleoside that is produced in tRNAs by the action of tRNA guanosine-2'-O-methyltransferase, using S-adenosyl-L-methionine as a substrate. Through its interaction with other modified nucleosides, 2'-O-methylguanosine is thought to stabilize the structure of the tRNA.

Recommended Dilutions

DB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot

Immunogen

Chemical compounds corresponding to 2'-O-Methylguanosine(Gm).

Synonyms

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
$\overline{\Box}$	Т	www.ahclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20 $^{\circ}\text{C}.$ Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



The 2'-O-methylguanosine(Gm) Rabbit pAb(A20697) are tested in Dot Blot against 2'-O-methylguanosine(Gm) and unmodified guanosine. Gm-3[5'Biotin-GCATAATGACTACT(Gm)3' Gm-4[5'Biotin-GCATAATGACTACTG3'