

# MYC2 Rabbit pAb

Catalog No.: A20774

## Basic Information

### Observed MW

75kDa

### Calculated MW

68kDa

### Category

Primary antibody

### Applications

WB, ELISA

### Cross-Reactivity

Arabidopsis thaliana, Other (Wide Range Predicted)

## Background

Encodes a MYC-related transcriptional activator with a typical DNA binding domain of a basic helix-loop-helix leucine zipper motif. Binds to an extended G-Box promoter motif and interacts with Jasmonate ZIM-domain proteins. Its transcription is induced by dehydration stress and ABA treatment. Negative regulator of blue light#Ae-mediated photomorphogenic growth and blue and far-red-light#Ae-regulated gene expression. Positive regulator of lateral root formation. Regulates diverse JA-dependent functions. Negatively regulates Trp metabolism and biosynthesis of Trp-derived secondary metabolites. Positively regulates flavonoid biosynthesis, resistance to insects, and response to oxidative stress. Regulates other transcription factors, and negatively regulates its own expression.

## Recommended Dilutions

**WB** 1:100 - 1:500

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

840158

### Swiss Prot

Q39204

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

ATMYC2; F6N18.4; F6N18\_4; JAI1; JASMONATE INSENSITIVE 1; JIN1; RD22BP1; ZBF1; MYC2

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.

# Validation Data



Western blot analysis of extracts of various tissues from *Arabidopsis thaliana*, using MYC2 Rabbit pAb (A20774) at 1:500 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 30s.