# **Bcl-2 Mouse mAb**

Catalog No.: A20777 12 Publications



## **Basic Information**

#### **Observed MW**

26 kDa

### **Calculated MW**

26 kDa

### Category

Primary antibody

## **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

#### CloneNo number

AMC0441

## **Background**

This gene encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Alternative splicing results in multiple transcript variants.

## **Recommended Dilutions**

WB 1:1000 - 1:3000

**IF/ICC** 1:50 - 1:200

 $\begin{array}{c} \textbf{ELISA} & \text{Recommended starting} \\ & \text{concentration is 1 } \mu\text{g/mL}. \end{array}$ 

Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**596

Swiss Prot
P10415

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

Bcl-2; PPP1R50

## **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\Box$	ī	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationMouseIgG1,KappaAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




Western blot analysis of various lysates using Bcl-2 Mouse mAb (A20777) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 10s.



Immunofluorescence analysis of HeLa cells using Bcl-2 Mouse mAb (A20777) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Mouse IgG (H+L) (AS008) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using Bcl-2 Mouse mAb (A20777) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Mouse IgG (H+L) (AS008) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using Bcl-2 Mouse mAb (A20777) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Mouse IgG (H+L) (AS008) at 1:500 dilution. Blue: DAPI for nuclear staining.