TNF-α Rabbit pAb

Catalog No.: A20851 6 Publications



Basic Information

Observed MW

25kDa/28kDa

Calculated MW

26kDa

Category

Primary antibody

Applications

ELISA,WB

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. Members of this family are classified based on primary sequence, function, and structure. This protein is synthesized as a type-II transmembrane protein and is reported to be cleaved into products that exert distinct biological functions. It plays an important role in the innate immune response as well as regulating homeostasis but is also implicated in diseases of chronic inflammation. In mouse deficiency of this gene is associated with defects in response to bacterial infection, with defects in forming organized follicular dendritic cell networks and germinal centers, and with a lack of primary B cell follicles. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB

1:500 - 1:1000

Immunogen Information

Gene ID 21926

Swiss Prot

P06804

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 80-235 of mouse TNF- α (NP_038721.1).

Synonyms

DIF; Tnfa; TNF-a; TNFSF2; Tnlg1f; Tnfsf1a; TNFalpha; TNF-alpha; TNF- α

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	T	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of lysates from RAW 264.7 cells using TNF- α Rabbit pAb (A20851) at 1:1000 dilution. Raw 264.7 cells were treated by LPS (1 μ g/ml) at 37°C for 8 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.