HNF-4-alpha Rabbit mAb

Catalog No.: A20865 Recombinant 6 Publications



Basic Information

Observed MW

53kDa

Calculated MW

53kDa

Category

Primary antibody

Applications

WB,IHC-P,ELISA,ChIP

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC2794

Background

The protein encoded by this gene is a transcription factor involved in the development of the pancreas, liver, kidney, and intestines. The encoded protein also functions to maintain glucose homeostasis. Several transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB 1:1000 - 1:6000

IHC-P 1:200 - 1:2000

ELISA Recommended starting

concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay

requirements.

ChIP 5μg antibody for

10μg-15μg of Chromatin

Immunogen Information

Gene ID Swiss Prot 15378 P49698

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Hnf4; HNF-4; MODY1; Nr2a1; Tcf14; TCF-14; Hnf4alpha; HNF-4-alpha

Contact

6	400-999-6126
\bowtie	cn.market@abclonal.com.cn
<u> </u>	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




Western blot analysis of various lysates using HNF-4-alpha Rabbit mAb (A20865) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 3s.




Chromatin immunoprecipitation analysis of extracts of HepG2 cells, using HNF-4-alpha antibody (A20865) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Immunohistochemistry analysis of paraffinembedded Human colon carcinoma using HNF-4-alpha Rabbit mAb (A20865) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human liver cancer using HNF-4-alpha Rabbit mAb (A20865) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human small intestine using HNF-4-alpha Rabbit mAb (A20865) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.