

Integrin alpha V (ITGAV/CD51) Rabbit pAb

Catalog No.: A2091 **4 Publications**

Basic Information

Observed MW

130kDa

Calculated MW

116kDa

Category

Primary antibody

Applications

WB,IHC-P,IP,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

The product of this gene belongs to the integrin alpha chain family. Integrins are heterodimeric integral membrane proteins composed of an alpha subunit and a beta subunit that function in cell surface adhesion and signaling. The encoded preproprotein is proteolytically processed to generate light and heavy chains that comprise the alpha V subunit. This subunit associates with beta 1, beta 3, beta 5, beta 6 and beta 8 subunits. The heterodimer consisting of alpha V and beta 3 subunits is also known as the vitronectin receptor. This integrin may regulate angiogenesis and cancer progression. Alternative splicing results in multiple transcript variants. Note that the integrin alpha 5 and integrin alpha V subunits are encoded by distinct genes.

Recommended Dilutions

WB 1:1000 - 1:5000

IHC-P 1:50 - 1:200

IP 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells

ELISA Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Immunogen Information

Gene ID

3685

Swiss Prot

P06756

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 950-1048 of human Integrin alpha V (ITGAV/CD51) (NP_002201.1).

Synonyms

CD51; MSK8; VNRA; VTNR; Integrin alpha V (ITGAV/CD51)

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

Validation Data



Immunoprecipitation analysis of 200 µg extracts of C6 cells using 3 µg Integrin alpha V (ITGAV/CD51) antibody (A2091). Western blot was performed from the immunoprecipitate using Integrin alpha V (ITGAV/CD51) antibody (A2091) at a dilution of 1:2000.



Western blot analysis of various lysates using Integrin alpha V (ITGAV/CD51) Rabbit pAb (A2091) at 1:3000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3s.



Western blot analysis of lysates from MCF7 cells, using Integrin alpha V (ITGAV/CD51) Rabbit pAb (A2091) at 1:3000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.

Validation Data



Immunohistochemistry analysis of paraffin-embedded Rat testis using Integrin alpha V (ITGAV/CD51) Rabbit pAb (A2091) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.