# **UCHL5IP Rabbit mAb**

Catalog No.: A21156 Recombinant



### **Basic Information**

#### **Observed MW**

40kDa

### **Calculated MW**

40kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Human

#### CloneNo number

ARC3021

# **Background**

This gene encodes a subunit of the augmin complex, which regulates centrosome and mitotic spindle integrity, and is necessary for the completion of cytokinesis. The encoded protein was identified by interaction with ubiquitin C-terminal hydrolase 37. Alternative splicing results in multiple transcript variants.

# **Recommended Dilutions**

**WB** 

1:500 - 1:1000

## **Immunogen Information**

**Gene ID** 55559

Swiss Prot

Q99871

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 150-285 of human UCHL5IP (NP $\_001372411.1$ ).

# **Synonyms**

UIP1; UCHL5IP

### **Contact**

<b>a</b>	400-999-6126
<b>×</b>	cn.market@abclonal.com.cn
$\overline{\Box}$	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of lysates from HeLa cells, using UCHL5IP Rabbit mAb (A21156) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit  $\lg G$  (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 120s.