# PGP9.5/UCHL1 Rabbit pAb

Catalog No.: A2131 2 Publications



### **Basic Information**

### **Observed MW**

27kDa

### **Calculated MW**

25kDa

### Category

Primary antibody

### **Applications**

ELISA,WB,IF/ICC

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The protein encoded by this gene belongs to the peptidase C12 family. This enzyme is a thiol protease that hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin. This gene is specifically expressed in the neurons and in cells of the diffuse neuroendocrine system. Mutations in this gene may be associated with Parkinson disease.

# **Recommended Dilutions**

**WB** 1:500 - 1:1000

**IF/ICC** 1:50 - 1:200

# **Immunogen Information**

**Gene ID**7345

Swiss Prot
P09936

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-220 of human PGP95/UCHL1 (NP\_004172.2).

# **Synonyms**

NDGOA; PARK5; PGP95; SPG79; PGP9.5; SPG79A; UCHL-1; Uch-L1; HEL-117; PGP 9.5; HEL-S-53; PGP9.5/UCHL1

## **Contact**

<b>a</b>	400-999-6126
×	cn.market@abclonal.com.cn
	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.




Western blot analysis of various lysates using PGP9.5/UCHL1 Rabbit pAb (A2131) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Immunofluorescence analysis of C6 cells using PGP9.5/PGP9.5/UCHL1 Rabbit pAb (A2131) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-251 MG cells using PGP9.5/PGP9.5/UCHL1 Rabbit pAb (A2131) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.