Peroxiredoxin 3 (PRDX3) Rabbit pAb

www.abclonal.com

ABclonal

Catalog No.: A21341

Basic Information

Observed MW

28kDa

Calculated MW

28kDa

Category

Primary antibody

Applications

ELISA,WB

Cross-Reactivity

Human, Rat

Background

This gene encodes a mitochondrial protein with antioxidant function. The protein is similar to the C22 subunit of Salmonella typhimurium alkylhydroperoxide reductase, and it can rescue bacterial resistance to alkylhydroperoxide in E. coli that lack the C22 subunit. The human and mouse genes are highly conserved, and they map to the regions syntenic between mouse and human chromosomes. Sequence comparisons with recently cloned mammalian homologs suggest that these genes consist of a family that is responsible for the regulation of cellular proliferation, differentiation and antioxidant functions. This family member can protect cells from oxidative stress, and it can promote cell survival in prostate cancer. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 1, 3, 13 and 22.

Recommended Dilutions

WB

1:1000 - 1:5000

Immunogen Information

Gene ID 10935

Swiss Prot

P30048

Immunogen

Recombinant Protein corresponding to a sequence within amino acids 63-256 of human Peroxiredoxin 3 (PRDX3) (NP_006784.1).

Synonyms

AOP1; MER5; AOP-1; PPPCD; SP-22; HBC189; SCAR32; PRO1748; prx-III; Peroxiredoxin 3 (PRDX3)

Contact

| a | | 400-999-6126 |
|-----------|---|---------------------------|
| \bowtie | | cn.market@abclonal.com.cn |
| • | Ī | www.abclonal.com.cn |

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20 $^{\circ}\text{C}.$ Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using Peroxiredoxin 3 (PRDX3) Rabbit pAb (A21341) at 1:3000 dilution

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates / proteins: 25 µg per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.