CD36/SR-B3 Rabbit pAb

Catalog No.: A21716



Basic Information

Observed MW

70-110kDa

Calculated MW

53kDa

Category

Primary antibody

Applications

ELISA,WB

Cross-Reactivity

Mouse, Rat

Background

The protein encoded by this gene is the fourth major glycoprotein of the platelet surface and serves as a receptor for thrombospondin in platelets and various cell lines. Since thrombospondins are widely distributed proteins involved in a variety of adhesive processes, this protein may have important functions as a cell adhesion molecule. It binds to collagen, thrombospondin, anionic phospholipids and oxidized LDL. It directly mediates cytoadherence of Plasmodium falciparum parasitized erythrocytes and it binds long chain fatty acids and may function in the transport and/or as a regulator of fatty acid transport. Mutations in this gene cause platelet glycoprotein deficiency. Multiple alternatively spliced transcript variants have been found for this gene.

Recommended Dilutions

WB

1:500 - 1:1000

Immunogen Information

Gene ID 948 Swiss Prot

P16671

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 93-165 of CD36/SR-B3 (NP_000063.2).

Synonyms

FAT; GP4; GP3B; GPIV; CHDS7; PASIV; SCARB3; BDPLT10; CD36/SR-B3

Contact

| <u>a</u> | | 400-999-6126 |
|-----------|---|---------------------------|
| \bowtie | | cn.market@abclonal.com.cn |
| \odot | 1 | www.abclonal.com.cn |


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates, using CD36/SR-B3 Rabbit pAb (A21716) at 1:800 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.