

[KO Validated] Ajuba Rabbit pAb

Catalog No.: A21760 **KO Validated**

Basic Information

Observed MW

55kDa/

Calculated MW

57kDa

Category

Primary antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Rat

Background

Enables alpha-catenin binding activity and transcription corepressor activity. Involved in several processes, including negative regulation of hippo signaling; positive regulation of gene silencing by miRNA; and regulation of cellular response to hypoxia. Acts upstream of or within gene silencing by miRNA and positive regulation of protein-containing complex assembly. Located in several cellular components, including Golgi apparatus; P-body; and nucleoplasm.

Recommended Dilutions

WB 1:500 - 1:1000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

84962

Swiss Prot

Q961F1

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

JUB; ba

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data



Western blot analysis of lysates from HepG2 cells, using [KO Validated] Ajuba Rabbit pAb (A21760) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90s.



Western blot analysis of lysates from Rat kidney using [KO Validated] Ajuba Rabbit pAb (A21760) at 1:1000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 180s.



Western blot analysis of lysates from wild type (WT) and AJUBA knockdown (KD) Hep G2 cells using [KO Validated] Ajuba Rabbit pAb (A21760) at 1:1000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90s.