# ABclonal www.abclonal.com

## ABflo® 488 Rabbit anti-Human CD13/ANPEP mAb

Catalog No.: A21944

#### **Basic Information**

#### **Observed MW**

**Calculated MW** 

110kDa

Category

Primary antibody

**Applications** 

FC

**Cross-Reactivity** 

Human

CloneNo number

ARC53703

Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

## **Background**

Aminopeptidase N is located in the small-intestinal and renal microvillar membrane, and also in other plasma membranes. In the small intestine aminopeptidase N plays a role in the final digestion of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases. Its function in proximal tubular epithelial cells and other cell types is less clear. The large extracellular carboxyterminal domain contains a pentapeptide consensus sequence characteristic of members of the zinc-binding metalloproteinase superfamily. Sequence comparisons with known enzymes of this class showed that CD13 and aminopeptidase N are identical. The latter enzyme was thought to be involved in the metabolism of regulatory peptides by diverse cell types, including small intestinal and renal tubular epithelial cells, macrophages, granulocytes, and synaptic membranes from the CNS. This membrane-bound zinc metalloprotease is known to serve as a receptor for the HCoV-229E alphacoronavirus as well as other non-human coronaviruses. This gene has also been shown to promote angiogenesis, tumor growth, and metastasis and defects in this gene are associated with various types of leukemia and lymphoma.

### **Recommended Dilutions**

FC

5  $\mu$ l per 10^6 cells in 100  $\mu$ l volume

## Immunogen Information

Gene ID

**Swiss Prot** 

P15144

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

#### **Synonyms**

APN; AP-M; AP-N; CD13; LAP1; P150; PEPN; hAPN; GP150

#### **Contact**

2		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

#### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## **Validation Data**



Flow cytometry: 1X10^6 Human PBMC were surface-stained with ABflo \$ 488 Rabbit IgG isotype control (A22069,5  $\mu$ l/Test,left) or ABflo \$ 488 Rabbit anti-Human CD13/ANPEP mAb (A21944,5  $\mu$ l/Test,right). Cells in the monocyte gate were used for analysis.