# **UCP1** Rabbit mAb

Catalog No.: A21979 Recombinant 1 Publications



## **Basic Information**

#### **Observed MW**

30kDa

#### **Calculated MW**

33kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Mouse

#### CloneNo number

ARC54141

# **Background**

Mitochondrial uncoupling proteins (UCP) are members of the family of mitochondrial anion carrier proteins (MACP). UCPs separate oxidative phosphorylation from ATP synthesis with energy dissipated as heat, also referred to as the mitochondrial proton leak. UCPs facilitate the transfer of anions from the inner to the outer mitochondrial membrane and the return transfer of protons from the outer to the inner mitochondrial membrane. They also reduce the mitochondrial membrane potential in mammalian cells. Tissue specificity occurs for the different UCPs and the exact methods of how UCPs transfer H+/OH- are not known. UCPs contain the three homologous protein domains of MACPs. This gene is expressed only in brown adipose tissue, a specialized tissue which functions to produce heat.

# **Recommended Dilutions**

**WB** 

1:500 - 1:1000

# **Immunogen Information**

**Gene ID** 7350

**Swiss Prot** 

P25874

### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 100-200 of human UCP1 ( $NP_068605.1$ ).

## **Synonyms**

UCP; SLC25A7; UCP1

### **Contact**

6		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.



Western blot analysis of various lysates, using UCP1 antibody (A21979) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.