# **NDUFAB1** Rabbit pAb

Catalog No.: A22019



### **Basic Information**

#### **Observed MW**

11kDa

### **Calculated MW**

17kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

Predicted to enable acyl binding activity; acyl carrier activity; and fatty acid binding activity. Involved in mitochondrial respiratory chain complex I assembly and protein lipoylation. Located in mitochondrion and nucleoplasm. Part of mitochondrial respiratory chain complex I. Colocalizes with mitochondrial large ribosomal subunit.

## **Recommended Dilutions**

WB

1:500 - 1:1000

## **Immunogen Information**

Gene ID 4706 **Swiss Prot** 

014561

### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 50-156 of human NDUFAB1 (NP\_004994.1).

### **Synonyms**

ACP; ACP1; SDAP; FASN2A; NDUFAB1

### **Contact**

|          | 400-999-6126              |
|----------|---------------------------|
| <b>×</b> | cn.market@abclonal.com.cn |
|          | www.abclonal.com.cn       |


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of extracts of various cell lines, using NDUFAB1 antibody (A22019) at 1:900 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG(H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.