DiMethyl-Histone H3-K36 Rabbit mAb



www.abclonal.com

Catalog No.: A22087 Recombinant 1 Publications

Basic Information

Observed MW

17 kDa

Calculated MW

15 kDa

Category

Primary antibody

Applications

WB, DB, IHC-P, IF/ICC, IP, ELISA

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

CloneNo number

ARC54175

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB 1:2000 - 1:10000 1:2000 - 1:10000 DB

IHC-P 1:1000 - 1:5000

1:50 - 1:200 IF/ICC

ΙP 2μg-6μg antibody for 300µg-600µg extracts of

whole cells

Recommended starting **ELISA**

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot 8290/8350 Q16695/P68431

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; DiMethyl-Histone H3-K36

Product Information

Source Isotype **Purification** Rabbit IgG Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cr
•	www.abclonal.com.cr



Immunoprecipitation analysis of 600 μ g extracts of 293F cells using 5 μ g DiMethyl-Histone H3-K36 antibody (A22087). Western blot was performed from the immunoprecipitate using DiMethyl-Histone H3-K36 antibody (A22087) at a dilution of 1:10000.



Western blot analysis of lysates from HeLa cells using DiMethyl-Histone H3-K36 Rabbit mAb (A22087) at 1:10000 dilution incubated overnight at 4° C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 μg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10 s.



Western blot analysis of various lysates using DiMethyl-Histone H3-K36 Rabbit mAb (A22087) at 1:10000 dilution incubated overnight at 4° C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).

Negative control (NC): Recombinant H3 protein(10 ng per lane).


Exposure time: 30 s.









Dot-blot analysis of all sorts of peptides using DiMethyl-Histone H3-K36 Rabbit mAb antibody (A22087) at 1:10000 dilution.



Immunohistochemistry analysis of paraffinembedded Human breast cancer tissue using DiMethyl-Histone H3-K36 Rabbit mAb (A22087) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human thyroid tissue using DiMethyl-Histone H3-K36 Rabbit mAb (A22087) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Rat liver tissue using DiMethyl-


Immunofluorescence analysis of C6 cells using DiMethyl-Histone H3-K36 Rabbit mAb (A22087) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffinembedded Human breast tissue using DiMethyl-Histone H3-K36 Rabbit mAb (A22087) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Mouse liver tissue using DiMethyl-Histone H3-K36 Rabbit mAb (A22087) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffinembedded Rat lung tissue using DiMethyl-

Immunofluorescence analysis of MCF7 cells using DiMethyl-Histone H3-K36 Rabbit mAb (A22087) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffinembedded Human esophagus tissue using DiMethyl-Histone H3-K36 Rabbit mAb (A22087) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Mouse spleen tissue using DiMethyl-Histone H3-K36 Rabbit mAb (A22087) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.

Validation Data

Histone H3-K36 Rabbit mAb (A22087) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.

Histone H3-K36 Rabbit mAb (A22087) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.