RGL2 Rabbit pAb

Catalog No.: A22119



Basic Information

Observed MW

60kDa

Calculated MW

60kDa

Category

Primary antibody

Applications

ELISA,WB

Cross-Reactivity

Arabidopsis thaliana

Background

Encodes a DELLA protein, a member of the GRAS superfamily of putative transcription factors. DELLA proteins restrain the cell proliferation and expansion that drives plant growth. Negative regulator of the response to GA in controlling seed germination. GA triggers the degradation of RGL2 protein in a process blocked by both proteasome inhibitors and serine/threonine phosphatase inhibitors. The protein undergoes degradation in response to GA via the 26S proteasome. RGL2 may be involved in reducing ROS accumulation in response to stress by up-regulating the transcription of superoxide dismutases. Rapidly degraded in response to GA. Regulates GA-promoted seed germination. Involved in flower and fruit development.

Recommended Dilutions

WB

1:1000 - 1:5000

Immunogen Information

Gene ID 821251

Swiss Prot

Q8GXW1

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 40-156 of arabidopsis thaliana RGL2 (NP_186995.1).

Synonyms

RGA-like 2; T21P5.13; T21P5_13; RGL2

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
\odot	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of various tissues from Arabidopsis thaliana, using RGL2 Rabbit pAb (A22119) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates / proteins: 25 µg per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021).

Exposure time: 60s.