ALKBH5 Rabbit mAb

Catalog No.: A22137 Recombinant



Basic Information

Observed MW

44-48kDa/45-48kDa

Calculated MW

44kDa

Category

Primary antibody

Applications

ELISA,WB,IHC-P,IP

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC55858

Background

Enables mRNA N6-methyladenosine dioxygenase activity. Involved in RNA metabolic process; mRNA export from nucleus; and response to hypoxia. Located in Golgi apparatus; cytosol; and nuclear speck.ALKBH5, also named as ABH5 and OFOXD1, is a Dioxygenase that demethylates RNA by oxidative demethylation. It is a mammalian demethylase that oxidatively reverses m6A in mRNA in vitro and in vivo. ALKBH5 is required for spermatogenesis.

Recommended Dilutions

WB 1:1000 - 1:5000

IHC-P 1:50 - 1:100

IP 0.5μg-4μg antibody for 400μg-600μg extracts of

whole cells

Immunogen Information

Gene IDSwiss Prot
54890
Q6P6C2

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 66-394 of human ALKBH5 (NP_060228.3).

Synonyms

ABH5; OFOXD; OFOXD1; ALKBH5

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	T	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.



Immunoprecipitation of ALKBH5 in 400 µg extracts from Rat brain using 0.5 µg ALKBH5 Rabbit mAb (A22137). Western blot analysis was performed using ALKBH5 Rabbit mAb (A22137) at 1:2000 dilution.




Western blot analysis of various lysates, using ALKBH5 Rabbit mAb (A22137) at 1:2000 dilution. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 30s.



Immunohistochemistry analysis of paraffinembedded human kidney using ALKBH5 Rabbit mAb (A22137) at dilution of 1:100 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffinembedded mouse brain using ALKBH5 Rabbit mAb (A22137) at dilution of 1:100 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffinembedded rat kidney using ALKBH5 Rabbit mAb (A22137) at dilution of 1:100 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.