

DiMethyl-Histone H3-K4 Rabbit mAb

Catalog No.: A22143

Recombinant

1 Publications

Basic Information

Observed MW

17kDa//17kDa

Calculated MW

16kDa

Category

Primary antibody

Applications

WB,DB,IHC-P,IF/ICC,IP,ELISA,ChIP,ChIP-seq,CUT&Tag

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

CloneNo number

ARC55489

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB 1:2000 - 1:10000**IP** 0.5µg-5µg antibody for
400µg-600µg extracts of
whole cells**IF/ICC** 1:50 - 1:200**IHC-P** 1:1000 - 1:5000**DB** 1:2000 - 1:6000**ChIP** 5µg antibody for
5µg-10µg of Chromatin**ChIP-seq** 1:50 - 1:200**CUT&Tag** 10⁵ cells /1 µg**ELISA** Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; DiMethyl-Histone H3-K4

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data




Immunoprecipitation of DiMethyl-Histone H3-K4 Rabbit from 600 µg extracts of HeLa cells was performed using 3 µg of DiMethyl-Histone H3-K4 Rabbit mAb (A22143). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1X Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using DiMethyl-Histone H3-K4 Rabbit mAb (A22143) at a dilution of 1:10000.



Immunoprecipitation of DiMethyl-Histone H3-K4 from 600 µg extracts of HeLa cells was performed using 3 µg of DiMethyl-Histone H3-K4 Rabbit mAb (A22143). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1X Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using DiMethyl-Histone H3-K4 Rabbit mAb (A22143) at a dilution of 1:10000.




Western blot analysis of lysates from C6 cells using DiMethyl-Histone H3-K4 Rabbit mAb (A22143) at 1:10000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.




Western blot analysis of various lysates using DiMethyl-Histone H3-K4 Rabbit mAb (A22143) at 1:10000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Negative control (NC): H3 protein
Exposure time: 20s.




Chromatin immunoprecipitation was performed with 10 µg of cross-linked chromatin from HeLa using 5 µg of DiMethyl-Histone H3-K4 Rabbit mAb [A22143]. DNA libraries were prepared using Scale ssDNA-seq Lib Prep Kit for Illumina V2 (RK20228). The ChIP sequencing results indicate the enrichment pattern of DiMethyl-Histone H3-K4 in the representative genomic region surrounding GAPDH gene.



Chromatin immunoprecipitation was performed with 10 μ g of cross-linked chromatin from HeLa using 5 μ g of DiMethyl-Histone H3-K4 Rabbit mAb (A22143). DNA libraries were prepared using Scale ssDNA-seq Lib Prep Kit for Illumina V2 (RK20228). The ChIP sequencing results indicate the enrichment pattern of DiMethyl-Histone H3-K4 across chromosome 12 (upper panel) and the genomic region encompassing GAPDH, a representative gene enriched in DiMethyl-Histone H3-K4 (lower panel).






Chromatin immunoprecipitation analysis of extracts of HeLa cells, using DiMethyl-Histone H3-K4 Rabbit mAb (A22295) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



CUT&Tag was performed using the CUT&Tag Assay Kit (pAG-Tn5) for Illumina (RK20265) from 10^5 HeLa with 1 μ g of DiMethyl-Histone H3-K4 Rabbit mAb (A22143), followed by incubation with Goat Anti-Rabbit IgG(H+L)(AS070). The CUT&Tag results denote the enrichment pattern of DiMethyl-Histone H3-K4 across chromosome 12 (upper panel) and the genomic region encompassing GAPDH, a representative gene enriched in DiMethyl-Histone H3-K4 (lower panel).


CUT&Tag was performed using the CUT&Tag Assay Kit (pAG-Tn5) for Illumina (RK20265) from 10⁵ HeLa cells with 1 µg of DiMethyl-Histone H3-K4 Rabbit mAb (A22143), followed by incubation with Goat Anti-Rabbit IgG(H+L)(AS070). The CUT&Tag results denote the enrichment pattern of DiMethyl-Histone H3-K4 around GAPDH gene.

Dot-blot analysis of all sorts of peptides using DiMethyl-Histone H3-K4 Rabbit mAb (A22143) at 1:5000 dilution.




Validation Data

Confocal imaging of HeLa cells using DiMethyl-Histone H3-K4 Rabbit mAb (A22143, dilution 1:200) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) (Green). DAPI was used for nuclear staining (blue). Objective: 100x.




Immunohistochemistry analysis of paraffin-embedded Human liver cancer tissue using DiMethyl-Histone H3-K4 Rabbit mAb (A22143) at a dilution of 1:1500 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.

Immunohistochemistry analysis of paraffin-embedded Human colon tissue using DiMethyl-Histone H3-K4 Rabbit mAb (A22143) at a dilution of 1:1500 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse colon tissue using DiMethyl-Histone H3-K4 Rabbit mAb (A22143) at a dilution of 1:1500 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.

Immunohistochemistry analysis of paraffin-embedded Human kidney tissue using DiMethyl-Histone H3-K4 Rabbit mAb (A22143) at a dilution of 1:1500 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat colon tissue using DiMethyl-Histone H3-K4 Rabbit mAb (A22143) at a dilution of 1:1500 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.