

Asymmetric DiMethyl-Histone H3-R8 Rabbit mAb

Catalog No.: A22144 **Recombinant**

Basic Information

Observed MW

17kDa

Calculated MW

16kDa

Category

Primary antibody

Applications

WB,DB,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

CloneNo number

ARC54909

Recommended Dilutions

WB 1:2000 - 1:4000**DB** 1:5000 - 1:8000**IHC-P** 1:50 - 1:200**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431

Immunogen

A synthetic asymmetric dimethylated peptide around R8 of human Histone H3 (NP_003520.1).

Synonyms

H3.4; H3/g; H3FT; H3t; HIST3H3; Histone H3; HIST1H3A; Asymmetric DiMethyl-Histone H3-R8

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.


Validation Data




Western blot analysis of various lysates, using Asymmetric DiMethyl-Histone H3-R8 Rabbit mAb (A22144) at 1:4000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 30s.




Dot-blot analysis of all sorts of peptides using Asymmetric DiMethyl-Histone H3-R8 Rabbit mAb (A22144) at 1:8200 dilution.




Immunohistochemistry analysis of paraffin-embedded Rat lung using Asymmetric DiMethyl-Histone H3-R8 Rabbit mAb (A22144) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse lung using Asymmetric DiMethyl-Histone H3-R8 Rabbit mAb (A22144) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon using Asymmetric DiMethyl-Histone H3-R8 Rabbit mAb (A22144) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human kidney using Asymmetric DiMethyl-Histone H3-R8 Rabbit mAb (A22144) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.