

NADPH oxidase 4 (NOX4) Rabbit mAb

Catalog No.: A22149 **Recombinant** **6 Publications**

Basic Information

Observed MW

62kDa

Calculated MW

67KDa

Category

Primary antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC55045

Background

This gene encodes a member of the NOX-family of enzymes that functions as the catalytic subunit the NADPH oxidase complex. The encoded protein is localized to non-phagocytic cells where it acts as an oxygen sensor and catalyzes the reduction of molecular oxygen to various reactive oxygen species (ROS). The ROS generated by this protein have been implicated in numerous biological functions including signal transduction, cell differentiation and tumor cell growth. A pseudogene has been identified on the other arm of chromosome 11. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB 1:1000 - 1:2000**IF/ICC** 1:100 - 1:400**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

50507

Swiss Prot

Q9NPH5

Immunogen

Recombinant protein of human NADPH oxidase 4 (NOX4)

Synonyms

KOX; KOX-1; RENOX; NADPH oxidase 4 (NOX4)

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates, using NADPH oxidase 4 (NOX4) Rabbit mAb (A22149) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Confocal imaging of HeLa cells using NADPH oxidase 4 (NOX4) Rabbit mAb (A22149, dilution 1:100) (Red). DAPI was used for nuclear staining (blue). Objective: 60x.