Leader in Biomolecular Solutions for Life Science



Catalog No.: A22197 Recombinant 2 Publications



Basic Information

Observed MW 90-120kDa

Calculated MW 41kDa

Category Primary antibody

Applications WB,IHC-P,ELISA

Cross-Reactivity Human

CloneNo number ARC55664

Background

The protein encoded by this gene may play a role in the attachment of stem cells to the bone marrow extracellular matrix or to stromal cells. This single-pass membrane protein is highly glycosylated and phosphorylated by protein kinase C. Two transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB	1:1000 - 1:5000
IHC-P	1:500 - 1:1000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID 947

Swiss Prot P28906

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 32-290 of human CD34 (NP_001020280.1).

Synonyms

CD34; CD34 molecule; GIG3; MORT1

Contact

2	400-999-6126
\mathbf{X}	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

Product Information


Source Rabbit

Isotype lgG


Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.



Western blot analysis of various lysates, using CD34 Rabbit mAb (A22197) at1:2000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:200000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 180s.



Immunohistochemistry analysis of paraffinembedded Human kidney tissue using CD34 Rabbit mAb (A22197) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.