

Acetyl-Histone H3-K27 Rabbit mAb

Catalog No.: A22264 **Recombinant** **3 Publications**

Basic Information

Observed MW

17kDa

Calculated MW

15kDa

Category

Primary antibody

Applications

ELISA,DB,WB,IHC-P,IF/ICC,ChIP,ChIP-seq

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

CloneNo number

ARC54943

Recommended Dilutions

DB 1:2000 - 1:20000

WB 1:2000 - 1:20000

IHC-P 1:50 - 1:200

IF/ICC 1:50 - 1:200

ChIP 5µg antibody for
5µg-10µg of Chromatin

ChIP-seq 1:50 - 1:200

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431

Immunogen

A synthetic acetylated peptide around K27 of human Histone H3 (NP_003520.1).

Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; Acetyl-Histone H3-K27

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.


Validation Data



Western blot analysis of lysates from HeLa cells, using Acetyl-Histone H3-K27 Rabbit mAb (A22264) at 1:20000 dilution. HeLa cells were treated by TSA (1 μ M) at 37°C for 18 hours.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 μ g per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.




Western blot analysis of lysates from NIH/3T3 cells, using Acetyl-Histone H3-K27 Rabbit mAb (A22264) at 1:20000 dilution. NIH/3T3 cells were treated by TSA (1 μ M) at 37°C for 18 hours.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 μ g per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.




Western blot analysis of lysates from C6 cells, using Acetyl-Histone H3-K27 Rabbit mAb (A22264) at 1:20000 dilution. C6 cells were treated by TSA (1 μ M) at 37°C for 18 hours.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 μ g per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.


Validation Data




Chromatin immunoprecipitations were performed with cross-linked chromatin from HeLa cells and Acetyl-Histone H3-K27 Rabbit mAb (A22264). The ChIP sequencing results indicate the enrichment pattern of Acetyl-Histone H3-K27 in selected genomic region and representative gene loci (RPL30), as shown in figure.




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using Acetyl-Histone H3-K27 antibody (A22264) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.




Dot-blot analysis of all sorts of peptides using Acetyl-Histone H3-K27 antibody (A22264) at 1:2000 dilution.




Immunohistochemistry analysis of paraffin-embedded Human lung using Acetyl-Histone H3-K27 Rabbit mAb (A22264) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human spleen using Acetyl-Histone H3-K27 Rabbit mAb (A22264) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-



Confocal imaging of HeLa TSA and HeLa cells

Validation Data

embedded Rat intestine using Acetyl-Histone H3-K27 Rabbit mAb (A22264) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.

using Acetyl-Histone H3-K27 Rabbit mAb (A22264,dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007,dilution 1:500)(Red).DAPI was used for nuclear staining (Blue). Objective: 100x.