

# TriMethyl-Histone H3-K27 Rabbit mAb

Catalog No.: A22396

Recombinant

6 Publications

## Basic Information

### Observed MW

17 kDa

### Calculated MW

15 kDa

### Category

Primary antibody

### Applications

WB, IF/ICC, IHC-P, ChIP, CUT&amp;Tag, ELISA

### Cross-Reactivity

Human, Mouse, Rat

### CloneNo number

ARC54167

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

## Recommended Dilutions

**WB** 1:20000 - 1:100000**IF/ICC** 1:1000 - 1:4000**IHC-P** 1:3000 - 1:12000**ChIP** 5 µg antibody for 5 µg-10 µg of Chromatin**CUT&Tag** 10<sup>5</sup> cells /1 µg**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

## Immunogen Information

### Gene ID

8290/8350

### Swiss Prot

Q16695/P68431

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; TriMethyl-Histone H3-K27

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




[www.abclonal.com.cn](http://www.abclonal.com.cn)

---


## Validation Data




Western blot analysis of various lysates using TriMethyl-Histone H3-K27 Rabbit mAb (A22396) at 1:40000 dilution incubated overnight at 4°C.  
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
 Lysates/proteins: 25 µg per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 45 s.




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using TriMethyl-Histone H3-K27 antibody (A22396) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



CUT&Tag was performed using the CUT&Tag Assay Kit (pAG-Tn5) for Illumina(RK20265) from 10<sup>5</sup> K562 cells with 1µg TriMethyl-Histone H3-K27 Rabbit mAb(A22396), along with a Goat Anti-Rabbit IgG(H+L). The CUT&Tag results indicate the enrichment pattern of H3K27Me3 in representative gene loci (MYT1), as shown in figure.



Immunofluorescence analysis of HeLa cells using TriMethyl-Histone H3-K27 Rabbit mAb (A22396) at dilution of 1:600 (40x lens).  
 Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded Human spleen using TriMethyl-Histone H3-K27 Rabbit mAb (A22396) at dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.