[KO Validated] MTA2 Rabbit pAb

www.abclonal.com

ABclonal

Catalog No.: A2243 KO Validated 1 Publications

Basic Information

Observed MW

75kDa

Calculated MW

75kDa

Category

Primary antibody

Applications

WB, ELISA, ChIP

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a protein that has been identified as a component of NuRD, a nucleosome remodeling deacetylase complex identified in the nucleus of human cells. It shows a very broad expression pattern and is strongly expressed in many tissues. It may represent one member of a small gene family that encode different but related proteins involved either directly or indirectly in transcriptional regulation. Their indirect effects on transcriptional regulation may include chromatin remodeling. It is closely related to another member of this family, a protein that has been correlated with the metastatic potential of certain carcinomas. These two proteins are so closely related that they share the same types of domains. These domains include two DNA binding domains, a dimerization domain, and a domain commonly found in proteins that methylate DNA. One of the proteins known to be a target protein for this gene product is p53. Deacetylation of p53 is correlated with a loss of growth inhibition in transformed cells supporting a connection between these gene family members and metastasis.

Recommended Dilutions

WB 1:500 - 1:1000

ELISA Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay

requirements.

5ug antibody for **ChIP**

10μg-15μg of Chromatin

Immunogen Information

Gene ID Swiss Prot 9219 094776

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

PID; MTA1L1; A2

Contact

a	400-999-6126
×	cn.market@abclonal.com.cn
$\overline{\alpha}$	www.ahclonal.com.cn


Product Information

Source Isotype **Purification** Rabbit Affinity purification IgG

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of lysates from wild type (WT) and MTA2 knockout (KO) 293T cells, using [KO Validated] MTA2 Rabbit pAb (A2243) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using MTA2 antibody (A2243) and rabbit IgG.The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.