

PI3 Kinase p85 alpha/beta Rabbit mAb

Catalog No.: A22487 **Recombinant** **1 Publications**

Basic Information

Observed MW

85kDa

Calculated MW

42-54kDa/81-84kDa

Category

Primary antibody

Applications

ELISA,WB

Cross-Reactivity

Human, Mouse

CloneNo number

ARC57210

Background

Phosphatidylinositol 3-kinase phosphorylates the inositol ring of phosphatidylinositol at the 3-prime position. The enzyme comprises a 110 kD catalytic subunit and a regulatory subunit of either 85, 55, or 50 kD. This gene encodes the 85 kD regulatory subunit.

Phosphatidylinositol 3-kinase plays an important role in the metabolic actions of insulin, and a mutation in this gene has been associated with insulin resistance. Alternative splicing of this gene results in four transcript variants encoding different isoforms. [provided by RefSeq, Jun 2011]

Recommended Dilutions

WB 1:500-1:1000

Immunogen Information

Gene ID

5295/5296

Swiss Prot

P27986/O00459

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 400-500 of human PI3 Kinase p85 alpha/beta (NP_852664.1).

Synonyms

PIK3R1/PIK3R2; PI3 Kinase p85 alpha/beta

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates, using PI3 Kinase p85 alpha/beta Rabbit mAb (A22487) at 1:3000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.