

Tamm Horsfall Glycoprotein Rabbit pAb

Catalog No.: A22544

Basic Information

Observed MW

105-120kDa

Calculated MW

70kDa

Category

Primary antibody

Applications

ELISA,WB,IF/ICC

Cross-Reactivity

Human, Mouse, Rat

Background

The protein encoded by this gene is the most abundant protein in mammalian urine under physiological conditions. Its excretion in urine follows proteolytic cleavage of the ectodomain of its glycosyl phosphatidylinositol-anchored counterpart that is situated on the luminal cell surface of the loop of Henle. This protein may act as a constitutive inhibitor of calcium crystallization in renal fluids. Excretion of this protein in urine may provide defense against urinary tract infections caused by uropathogenic bacteria. Defects in this gene are associated with the renal disorders medullary cystic kidney disease-2 (MCKD2), glomerulocystic kidney disease with hyperuricemia and isosthenuria (GCKDHI), and familial juvenile hyperuricemic nephropathy (FJHN). Alternative splicing of this gene results in multiple transcript variants.

Recommended Dilutions

WB	1:100 - 1:500
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

7369

Swiss Prot

P07911

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 175-324 of human Tamm Horsfall Glycoprotein. (NP_003352.2).

Synonyms

THP; FJHN; HNFJ; THGP; HNFJ1; MCKD2; ADTKD1; ADMCKD2; Tamm Horsfall Glycoprotein

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Western blot analysis of Mouse kidney, using Tamm Horsfall Glycoprotein antibody (A22544) at 1:400 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 60s.



Immunofluorescence analysis of Human kidney using Tamm Horsfall Glycoprotein Rabbit pAb (A22544) at dilution of 1:20 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of Mouse kidney using Tamm Horsfall Glycoprotein Rabbit pAb (A22544) at dilution of 1:20 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of Rat Kidney using Tamm Horsfall Glycoprotein Rabbit pAb (A22544) at dilution of 1:20 (40x lens). Blue: DAPI for nuclear staining.