ABflo® 647 Rabbit anti-Dog CD27 mAb

ABclonal www.abclonal.com

Catalog No.: A22576

Basic Information

Observed MW

Calculated MW

31kDa

Category

Primary antibody

Applications

Cross-Reactivity

Dog

CloneNo number

ARC55676

Conjugate

ABflo® 647, Ex:648nm, Em:664nm,

Background

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor.

Recommended Dilutions

FC

5 µl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID

Swiss Prot 611674

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

T14; S152; Tp55; TNFRSF7; S152. LPFS2

Contact

2		400-999-6126
\bowtie	Τ	cn.market@abclonal.com.cn
•	Т	www.abclonal.com.cn

Product Information

Source Isotype **Purification** lgG Rabbit Affinity purification

Storage


Store at 2-8°C. Avoid freeze.


Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data









Flow cytometry:1X10^6 293F cells (negative control,left) and 293F(Transfection,right) cells were surface-stained with ABflo® 647 Rabbit anti-Dog CD27 mAb(A22576,5 µl/Test,orange line) or ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:1X10^6 293F(Transfection) cells were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,left) or ABflo® 647 Rabbit anti- Dog CD27 mAb(A22576,5 µl/Test,right).

Flow cytometry: 1X106 Dog splenocytes were surface-stained with ABflo & 647 Rabbit IgG isotype control (A22070,5 μ l/Test,left) or ABflo & 647 Rabbit anti-Dog CD27 mAb (A22576,5 μ l/Test,right). Cells in the lymphocyte gate were used for analysis.