# ABflo® 488 Rabbit anti-Human CD304/Neuropilin-1 mAb

Catalog No.: A22579



## **Basic Information**

#### **Observed MW**

Calculated MW 68kDa/71kDa/103kDa

Category Primary antibody

Applications FC

Cross-Reactivity Human

CloneNo number ARC55418-ABf488

### Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

## **Recommended Dilutions**

FC

5 μl per 10^6 cells in 100 μl volume

# Background

This gene encodes one of two neuropilins, which contain specific protein domains which allow them to participate in several different types of signaling pathways that control cell migration. Neuropilins contain a large N-terminal extracellular domain, made up of complement-binding, coagulation factor V/VIII, and meprin domains. These proteins also contains a short membrane-spanning domain and a small cytoplasmic domain. Neuropilins bind many ligands and various types of co-receptors; they affect cell survival, migration, and attraction. Some of the ligands and co-receptors bound by neuropilins are vascular endothelial growth factor (VEGF) and semaphorin family members. This protein has also been determined to act as a co-receptor for SARS-CoV-2 (which causes COVID-19) to infect host cells.

## Immunogen Information

Gene ID 8829 **Swiss Prot** 014786

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 22-644 of human CD304/Neuropilin-1 (NP\_003864.5).

### Synonyms

NP1; NRP; BDCA4; CD304; VEGF165R

## Contact

6	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
€	www.abclonal.com.cn


## **Product Information**

**Source** Rabbit **Isotype** IgG Purification Affinity purification

## Storage

Store at 2-8°C. Avoid freeze. Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

## Validation Data







Flow cytometry:1X10^6 K-562 cells (negative control,left) and A549 cells (right) were surface-stained with ABflo® 488 Rabbit anti-Human CD304/Neuropilin-1 mAb(A22579,5 µl/Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line). Flow cytometry:1X10^6 A549 cells were surface-stained with ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,left) or ABflo® 488 Rabbit anti-Human CD304/Neuropilin-1 mAb(A22579,5 µl/Test,right).