

DNMT3B Rabbit mAb

Catalog No.: A22658 **Recombinant**

Basic Information

Observed MW

100kDa

Calculated MW

96kDa

Category

Primary antibody

Applications

WB, ELISA, ChIP, ChIP-seq

Cross-Reactivity

Human, Mouse

CloneNo number

ARC57702

Background

CpG methylation is an epigenetic modification that is important for embryonic development, imprinting, and X-chromosome inactivation. Studies in mice have demonstrated that DNA methylation is required for mammalian development. This gene encodes a DNA methyltransferase which is thought to function in de novo methylation, rather than maintenance methylation. The protein localizes primarily to the nucleus and its expression is developmentally regulated. Mutations in this gene cause the immunodeficiency-centromeric instability-facial anomalies (ICF) syndrome. Eight alternatively spliced transcript variants have been described. The full length sequences of variants 4 and 5 have not been determined.

Recommended Dilutions

WB 1:1000 - 1:5000**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.**ChIP** 5µg antibody for 10µg-15µg of Chromatin**ChIP-seq** 1:50 - 1:100

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Immunogen Information

Gene ID

1789

Swiss Prot

Q9UBC3

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

ICF; ICF1; FSHD4; M.HsaIIIB; DNMT3B

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of various lysates using DNMT3B Rabbit mAb (A22658) at 1:2000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 60s.



Chromatin immunoprecipitation was performed with 14 µg of cross-linked chromatin from A-549 cells using 5 µg of DNMT3B Rabbit mAb (A22658). DNA libraries were prepared using Scale ssDNA-seq Lib Prep Kit for Illumina V2 (RK20228). The ChIP sequencing results indicate the enrichment pattern of DNMT3B across chromosome X (upper panel) and the genomic region encompassing HUWE1, a representative gene enriched in DNMT3B (lower panel).



Chromatin immunoprecipitation was performed with 14 µg of cross-linked chromatin from NCCIT cells using 5 µg of DNMT3B Rabbit mAb (A22658). DNA libraries were prepared using Scale ssDNA-seq Lib Prep Kit for Illumina V2 (RK20228). The ChIP sequencing results indicate the enrichment pattern of DNMT3B in the representative genomic region surrounding HUWE1 gene.



Chromatin immunoprecipitation was performed with 10 µg of cross-linked chromatin from NCCIT cells, using 5 µg of DNMT3B antibody (A22658) and Rabbit IgG isotype control (AC042). The enrichment of immunoprecipitated DNA at different genomic loci was examined by quantitative PCR. The histogram compares the ratio of the immunoprecipitated DNA to the input at given loci.