

ABflo® 594 Rabbit anti-Goat IgG (Fab) fragment mAb

Catalog No.: A22785 **1 Publications**

Basic Information

Observed MW

Calculated MW

Category

Primary antibody

Applications

FC

Cross-Reactivity

Goat

CloneNo number

ARC57078-ABf594

Conjugate

ABflo® 594. Ex:588nm. Em:604nm.

Background

Recommended Dilutions

FC 5 µl per 10⁶ cells in
100 µl volume

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Recombinant protein of Goat IgG (Fab) fragment.

Synonyms

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data



Flow cytometry: 1×10^6 Jurkat cells (negative control, left) and A-431 cells (right) were surface-stained with Goat anti-Human TROP-2 mAb (5 μ l/Test, orange line) or ABflo® 594 Rabbit IgG isotype control (A23821, 5 μ l/Test, blue line), then stained with ABflo® 594 Rabbit anti-Goat IgG (Fab) fragment mAb (A22785, 5 μ l/Test) was used as a secondary antibody. Non-fluorescently stained Jurkat cells were used as blank control (red line).

Flow cytometry: 1×10^6 A-431 cells were surface-stained with ABflo® 594 Rabbit IgG isotype control (A23821, 5 μ l/Test, left) or Goat anti-Human TROP-2 mAb (5 μ l/Test, orange line). ABflo® 594 Rabbit anti-Goat IgG (Fab) fragment mAb (A22785, 5 μ l/Test) was used as a secondary antibody.