Leader in Biomolecular Solutions for Life Science

www.abclonal.com

PE/Cyanine7 Mouse anti-Human CD19 mAb

Catalog No.: A22819

Basic Information

Observed MW
Refer to figures

Calculated MW
61kDa

Category
Primary antibody

Applications
FC

Cross-Reactivity
Human

CloneNo number
AMC0541

Conjugate
PE-Cy7. Ex:565nm. Em:778nm.

Background

This gene encodes a member of the immunoglobulin gene superfamily. Expression of this cell surface protein is restricted to B cell lymphocytes. This protein is a reliable marker for pre-B cells but its expression diminishes during terminal B cell differentiation in antibody secreting plasma cells. The protein has two N-terminal extracellular Ig-like domains separated by a non-Ig-like domain, a hydrophobic transmembrane domain, and a large C-terminal cytoplasmic domain. This protein forms a complex with several membrane proteins including complement receptor type 2 (CD21) and tetraspanin (CD81) and this complex reduces the threshold for antigen-initiated B cell activation. Activation of this B-cell antigen receptor complex activates the phosphatidylinositol 3-kinase signalling pathway and the subsequent release of intracellular stores of calcium ions. This protein is a target of chimeric antigen receptor (CAR) T-cells used in the treatment of lymphoblastic leukemia. Mutations in this gene are associated with the disease common variable immunodeficiency 3 (CVID3) which results in a failure of B-cell differentiation and impaired secretion of immunoglobulins. CVID3 is characterized by hypogammaglobulinemia, an inability to mount an antibody response to antigen, and recurrent bacterial infections. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Recommended Dilutions

FC	10 µL/Test

Immunogen Information

Gene ID
930

Swiss Prot
P15391

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
B4; CVID3

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information


Source
Mouse

Isotype
IgG1


Purification
Affinity purification

Storage
Store at -20℃

Validation Data



Flow cytometry:1X10^6 Human PBMC were surface-stained with PE-Cy7 Mouse anti-Human CD19 mAb (A22819, orange line) or isotype control mAb (blue line).



Flow cytometry:1X10^6 Human PBMC were surface-stained with Isotype control mAb (left) or PE-Cy7 Mouse anti-Human CD19 mAb (A22819, right).