

# Acetyl-Histone H4-K5 Rabbit mAb

Catalog No.: A23080 **Recombinant**

## Basic Information

### Observed MW

11kDa

### Calculated MW

11kDa

### Category

Primary antibody

### Applications

WB, IHC-P, ELISA, ChIP, ChIP-seq, CUT&amp;Tag

### Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

### CloneNo number

ARC58041

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6.

## Recommended Dilutions

**WB** 1:2000 - 1:10000**IHC-P** 1:500 - 1:1000**ChIP** 5µg antibody for  
5µg-10µg of Chromatin**ChIP-seq** 1:50 - 1:200**CUT&Tag** 10<sup>5</sup> cells / 1 µg**ELISA** Recommended starting  
concentration is 1 µg/mL.  
Please optimize the  
concentration based on  
your specific assay  
requirements.

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

## Immunogen Information

### Gene ID

8359

### Swiss Prot

P62805

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

H4C2; H4C3; H4C4; H4C5; H4C6; H4C8; H4C9; H4FA; H4-16; H4C11; H4C12; H4C13; H4C14; H4C15; H4C16; HIST1H4A; Acetyl-Histone H4-K5

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




[www.abclonal.com.cn](http://www.abclonal.com.cn)

---


Validation Data



Western blot analysis of lysates from HeLa cells, using Acetyl-Histone H4-K5 Rabbit mAb (A23080) at 1:10000 dilution. HeLa cells were treated with TSA (1 uM) at 37°C for 18 hours.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 30s.





Western blot analysis of lysates from C2C12 cells, using Acetyl-Histone H4-K5 Rabbit mAb (A23080) at 1:10000 dilution. C2C12 cells were treated with TSA (1 uM) at 37°C for 18 hours.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 30s.




Western blot analysis of lysates from C6 cells, using Acetyl-Histone H4-K5 Rabbit mAb (A23080) at 1:10000 dilution. C6 cells were treated with TSA (1 uM) at 37°C for 18 hours.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 30s.

Validation Data


Chromatin immunoprecipitations were performed with cross-linked chromatin from HeLa cells and Acetyl-Histone H4-K5 Rabbit mAb (A23080). The ChIP sequencing results indicate the enrichment pattern of Acetyl-Histone H4-K5 in selected genomic region and representative gene loci (GAPDH), as shown in figure.




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using Acetyl-Histone H4-K5 antibody (A23080) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.




Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using Acetyl-Histone H4-K5 Rabbit mAb (A23080) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse intestine tissue using Acetyl-Histone H4-K5 Rabbit mAb (A23080) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse kidney tissue using Acetyl-Histone H4-K5 Rabbit mAb (A23080) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse liver tissue using Acetyl-Histone H4-K5 Rabbit mAb (A23080) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat intestine tissue using Acetyl-Histone H4-K5 Rabbit mAb (A23080) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat liver tissue using Acetyl-Histone H4-K5 Rabbit mAb (A23080) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



CUT&Tag was performed using the CUT&Tag Assay Kit (pAG-Tn5) for Illumina(RK20265) from 10<sup>5</sup> K562 cells with 1 μg Acetyl-Histone H4-K5 Rabbit mAb (A23080), along with a Goat Anti-Rabbit IgG(H+L). The CUT&Tag results indicate the enrichment pattern of Acetyl-Histone H4-K5 in representative gene loci (GAPDH), as shown in figure.