# ABflo® 647 Rabbit anti-Human B7-H2/CD275 mAb

Background

Catalog No.: A23104



## **Basic Information**

**Observed MW** 

Calculated MW 20kDa/33kDa34kDa

Category Primary antibody

Applications FC

Cross-Reactivity Human

CloneNo number ARC58551-ABf647

**Conjugate** ABflo® 647. Ex:648nm. Em:664nm.

## **Recommended Dilutions**

5 µl per 10^6 cells in

100 µl volume

FC

## Immunogen Information

**Gene ID** 23308

**Swiss Prot** 075144

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 19-256 of human B7-H2/CD275 (NP\_056074.1).

Enables identical protein binding activity. Predicted to be involved in T cell receptor signaling

pathway and positive regulation of interleukin-4 production. Located in cytoplasmic

ribonucleoprotein granule and plasma membrane.

#### Synonyms

B7h; B7H2; GL50; B7-H2; B7RP1; CD275; ICOSL; LICOS; B7RP-1; ICOS-L

## Contact

6	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
€	www.abclonal.com.cn


## **Product Information**


**Source** Rabbit **lsotype** IgG **Purification** Affinity purification

#### Storage

Store at 2-8°C. Avoid freeze. Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

## Validation Data







Flow cytometry:1X10^6 SH-SY5Y cells (negative control,Left) and Daudi cells (Right) were surface-stained with ABflo® 647 Rabbit anti-Human B7-H2/CD275 mAb(A23104,5 µl/Test,orange line) or ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,blue line). Non-fluorescently stained cells was used as blank control (red line). Flow cytometry:1X10<sup>6</sup> Daudi cells were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,left) or ABflo® 647 Rabbit anti-Human B7-H2/CD275 mAb(A23104,5 µl/Test,right).