

# ABflo® 488 Rabbit anti-Mouse IgM mAb

**Catalog No.: A23114**

## Basic Information

### Observed MW

Refer to figures

### Calculated MW

50kDa

### Category

Primary antibody

### Applications

FC

### Cross-Reactivity

Mouse

### CloneNo number

ARC58894

### Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

## Recommended Dilutions

**FC** 5  $\mu$ l per  $10^6$  cells in 100  $\mu$ l volume

## Background

Enables several functions, including antigen binding activity; identical protein binding activity; and immunoglobulin receptor binding activity. Acts upstream of or within several processes, including early endosome to late endosome transport; immunoglobulin mediated immune response; and positive regulation of B cell proliferation. Located in external side of plasma membrane; extracellular space; and perinuclear region of cytoplasm. Part of B cell receptor complex and immunoglobulin complex, circulating. Is expressed in several structures, including brain; hemolymphoid system; and liver. Used to study non-Hodgkin lymphoma and type 1 diabetes mellitus. Human ortholog(s) of this gene implicated in agammaglobulinemia. Orthologous to human IGHM (immunoglobulin heavy constant mu).

## Immunogen Information

### Gene ID

**Swiss Prot**  
P01872

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

Igm; muH; Igh6; Igh-6; Igh-M

## Contact

	400-999-6126
	<a href="mailto:cn.market@abclonal.com.cn">cn.market@abclonal.com.cn</a>
	<a href="http://www.abclonal.com.cn">www.abclonal.com.cn</a>

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


## Validation Data




Flow cytometry:  $1 \times 10^6$  NIH/3T3 cells (negative control, left) and C57BL/6 mouse splenocytes (right) were surface-stained with ABflo® 488 Rabbit anti-Mouse IgM mAb (A23114,5  $\mu$ l/Test, orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5  $\mu$ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry:  $1 \times 10^6$  C57BL/6 mouse splenocytes were surface-stained with ABflo® 488 Rabbit IgG isotype control (A22069,5  $\mu$ l/Test, left) or ABflo® 488 Rabbit anti-Mouse IgM mAb (A23114,5  $\mu$ l/Test, right).



Flow cytometry:  $1 \times 10^6$  NIH/3T3 cells (negative control, left) and Mouse PBMC (right) were surface-stained with ABflo® 488 Rabbit anti-Mouse IgM mAb (A23114,5  $\mu$ l/Test, orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5  $\mu$ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry:  $1 \times 10^6$  Mouse PBMC were surface-stained with ABflo® 488 Rabbit IgG isotype control (A22069,5  $\mu$ l/Test, left) or ABflo® 488 Rabbit anti-Mouse IgM mAb (A23114,5  $\mu$ l/Test, right).